Author Index Vol. 691

Acworth, I.N., see McCabe, D.R. 691(1997)23
Anspach, F.B., see Legallais, C. 691(1997)33
Asmis, R., Bühler, E., Jelk, J. and Gey, K.F.
Concurrent quantification of cellular cholesterol, cholesteryl esters and triglycerides in small biological samples. Reevaluation of thin layer chromatography using laser densitometry 691(1997)59
Atassi, G., see Tassin, J.P. 691(1997)449
Azuma, R. and Urakawa, A.
Simultaneous determination of a novel anticancer drug, TAS-103, and its N-demethylated metabolite in monkey plasma by high-performance liquid chromatography using solid-phase extraction 691(1997)179
Berger, B.J., Suskin, M., Dai, W.W., Cerami, A. and Ulrich, P.
Studies on the pharmacological properties of novel arylene bis(methylketone) compounds using solid-phase extraction and high-performance liquid chromatography 691(1997)433
Blaas, D., see Morand, M. 691(1997)192
Blaschke, G., see Paris, S. 691(1997)463
Blomhoff, R., see Gundersen, Th.E. 691(1997)43
Borbe, H.O., see Paris, S. 691(1997)463
Boudinot, F.D., see Chen, H. 691(1997)425
Brekelmans, G.J.F., see Saris, L.A. 691(1997)409
Bressolle, F., see Marquer, C. 691(1997)389
Brooks, S.A., Lachno, D.R. and Obermeyer, B.D.
Automated high-performance liquid chromatographic method for the analysis of two novel ergoline compounds in human plasma 691(1997)383
Brückner, H., see Schieber, A. 691(1997)1
Brulez, H.F.H., see Teerlink, T. 691(1997)269
Bueno, S.M.A., see Legallais, C. 691(1997)33
Bühler, E., see Asmis, R. 691(1997)59
Burke, T.G., see Warner, D.L. 691(1997)161
Cai, J. and Henion, J.
Quantitative multi-residue determination of β-agonists in bovine urine using on-line immunoaffinity extraction–coupled column packed capillary liquid chromatography–tandem mass spectrometry 691(1997)357
Cerami, A., see Berger, B.J. 691(1997)433
Chavez-Eng, C.M., Constanzer, M.L. and Matuszewski, B.K.
Picogram determination of a novel dopamine D_4 receptor antagonist in human plasma and urine by liquid chromatography with atmospheric pressure chemical ionization tandem mass spectrometry 691(1997)77
Chen, H., Manouilov, K.K., Chu, C.K., Schinazi, R.F., McClure, H.M. and Boudinot, F.D.
High-performance liquid chromatographic determination of (−)-β-D-2-aminopurine dioxolane and (−)-β-D-2-amino-6-chloropurine dioxolane, and their metabolite (−)-β-D-dioxolane guanine in monkey serum, urine and cerebrospinal fluid 691(1997)425
Chiu, F.C.K., Zhang, J.N., Li, R.C. and Raymond, K.
Efficient assay for the determination of atenolol in human plasma and urine by high-performance liquid chromatography with fluorescence detection 691(1997)473
Chu, C.K., see Chen, H. 691(1997)425
Classen, H.-G., see Schieber, A. 691(1997)1
Coleman, N.L., see Farid, N.A. 691(1997)212
Constanzer, M.L., see Chavez-Eng, C.M. 691(1997)77
Dai, W.W., see Berger, B.J. 691(1997)433
Davies, N.M.
Methods of analysis of chiral non-steroidal anti-inflammatory drugs 691(1997)229
Desiderio, D.M., see Lee, H.G. 691(1997)67
Deutz, N.E.P., see Van Eijk, H.M.H. 691(1997)287
Donker, M., see Sandrenan, N. 691(1997)145
Drewe, J., Rufer, S., Huwyler, J. and Küsters, E.
High-performance liquid chromatographic method for an automated determination of local anaesthetics in human plasma 691(1997)105
Dubois, J., see Tassin, J.P. 691(1997)449
Edelbroek, P.M., see Saris, L.A. 691(1997)409
El-Yazigi, A. and Legayada, E.
Direct liquid chromatographic micro-measurement of tamoxifen in plasma of cancer patients 691(1997)457
Engel, J., see Paris, S. 691(1997)463
Erdin, R., see Timm, U. 691(1997)397
Farid, N.A. and Coleman, N.L.
High-performance liquid chromatographic determination of the enantiomers of the benzoquinolinone LY191704, a human type 1 5α-reductase inhibitor, in plasma 691(1997)212
Feng, Y., Halaris, A.E. and Piletz, J.E.
Determination of agmatine in brain and plasma using high-performance liquid chromatography with fluorescence detection 691(1997)277
Gage, R., see Stopher, D.A. 691(1997)441
Gao, Q., see Manabe, S. 691(1997)197
Gey, K.F., see Asmis, R. 691(1997)59
Godbillon, J., see Sandrenan, N. 691(1997)145
Gundersen, Th.E., Lundanes, E. and Blomhoff, R.
Quantitative high-performance liquid chromatographic determination of retinoids in human serum using on-line solid-phase extraction and column switching. Determination of 9-*cis*-retinoic acid, 13-*cis*-retinoic acid, all-*trans*-retinoic acid, 4-oxo-all-*trans*-retinoic acid and 4-oxo-13-*cis*-retinoic acid 691(1997)43
Halaris, A.E., see Feng, Y. 691(1997)277
Hampl, R., see Hill, M. 691(1997)187
Hanocq, M., see Tassin, J.P. 691(1997)449
Haupt, K., see Legallais, C. 691(1997)33
Heimbecher, S., Lee, Y.-C., Tabibi, S.E. and Yalkowsky, S.H.
Derivatization and high-performance liquid chromatographic analysis of pentaazapentacosane pentahydrochloride 691(1997)173
Hemminki, K., see Koskinen, M. 691(1997)155
Henion, J., see Cai, J. 691(1997)357
Hennekes, M.W.T., see Teerlink, T. 691(1997)269
Hibiya, M., Teradaira, R., Shimpo, K., Matsui, T., Sugimoto, T. and Nagatsu, T.
Interference of a methotrexate derivative with urinary oncopterin [N^2-(3-aminopropyl)biopterin] measurement by high-performance liquid chromatography with fluorimetric detection 691(1997)223

Author Index Vol. 691

Acworth, I.N., see McCabe, D.R. 691(1997)23
Anspach, F.B., see Legallais, C. 691(1997)33
Asmis, R., Bühler, E., Jelk, J. and Gey, K.F.
Concurrent quantification of cellular cholesterol, cholesteryl esters and triglycerides in small biological samples. Reevaluation of thin layer chromatography using laser densitometry 691(1997)59
Atassi, G., see Tassin, J.P. 691(1997)449
Azuma, R. and Urakawa, A.
Simultaneous determination of a novel anticancer drug, TAS-103, and its N-demethylated metabolite in monkey plasma by high-performance liquid chromatography using solid-phase extraction 691(1997)179
Berger, B.J., Suskin, M., Dai, W.W., Cerami, A. and Ulrich, P.
Studies on the pharmacological properties of novel arylene bis(methylketone) compounds using solid-phase extraction and high-performance liquid chromatography 691(1997)433
Blaas, D., see Morand, M. 691(1997)192
Blaschke, G., see Paris, S. 691(1997)463
Blomhoff, R., see Gundersen, Th.E. 691(1997)43
Borbe, H.O., see Paris, S. 691(1997)463
Boudinot, F.D., see Chen, H. 691(1997)425
Brekelmans, G.J.F., see Saris, L.A. 691(1997)409
Bressolle, F., see Marquer, C. 691(1997)389
Brooks, S.A., Lachno, D.R. and Obermeyer, B.D.
Automated high-performance liquid chromatographic method for the analysis of two novel ergoline compounds in human plasma 691(1997)383
Brückner, H., see Schieber, A. 691(1997)1
Brulez, H.F.H., see Teerlink, T. 691(1997)269
Bueno, S.M.A., see Legallais, C. 691(1997)33
Bühler, E., see Asmis, R. 691(1997)59
Burke, T.G., see Warner, D.L. 691(1997)161
Cai, J. and Henion, J.
Quantitative multi-residue determination of β-agonists in bovine urine using on-line immunoaffinity extraction–coupled column packed capillary liquid chromatography–tandem mass spectrometry 691(1997)357
Cerami, A., see Berger, B.J. 691(1997)433
Chavez-Eng, C.M., Constanzer, M.L. and Matuszewski, B.K.
Picogram determination of a novel dopamine D_4 receptor antagonist in human plasma and urine by liquid chromatography with atmospheric pressure chemical ionization tandem mass spectrometry 691(1997)77
Chen, H., Manouilov, K.K., Chu, C.K., Schinazi, R.F., McClure, H.M. and Boudinot, F.D.
High-performance liquid chromatographic determination of (−)-β-D-2-aminopurine dioxolane and (−)-β-D-2-amino-6-chloropurine dioxolane, and their metabolite (−)-β-D-dioxolane guanine in monkey serum, urine and cerebrospinal fluid 691(1997)425
Chiu, F.C.K., Zhang, J.N., Li, R.C. and Raymond, K.
Efficient assay for the determination of atenolol in human plasma and urine by high-performance liquid chromatography with fluorescence detection 691(1997)473
Chu, C.K., see Chen, H. 691(1997)425
Classen, H.-G., see Schieber, A. 691(1997)1
Coleman, N.L., see Farid, N.A. 691(1997)212
Constanzer, M.L., see Chavez-Eng, C.M. 691(1997)77
Dai, W.W., see Berger, B.J. 691(1997)433
Davies, N.M.
Methods of analysis of chiral non-steroidal anti-inflammatory drugs 691(1997)229
Desiderio, D.M., see Lee, H.G. 691(1997)67
Deutz, N.E.P., see Van Eijk, H.M.H. 691(1997)287
Donker, M., see Sandrenan, N. 691(1997)145
Drewe, J., Rufer, S., Huwyler, J. and Küsters, E.
High-performance liquid chromatographic method for an automated determination of local anaesthetics in human plasma 691(1997)105
Dubois, J., see Tassin, J.P. 691(1997)449
Edelbroek, P.M., see Saris, L.A. 691(1997)409
El-Yazigi, A. and Legayada, E.
Direct liquid chromatographic micro-measurement of tamoxifen in plasma of cancer patients 691(1997)457
Engel, J., see Paris, S. 691(1997)463
Erdin, R., see Timm, U. 691(1997)397
Farid, N.A. and Coleman, N.L.
High-performance liquid chromatographic determination of the enantiomers of the benzoquinolinone LY191704, a human type 1 5α-reductase inhibitor, in plasma 691(1997)212
Feng, Y., Halaris, A.E. and Piletz, J.E.
Determination of agmatine in brain and plasma using high-performance liquid chromatography with fluorescence detection 691(1997)277
Gage, R., see Stopher, D.A. 691(1997)441
Gao, Q., see Manabe, S. 691(1997)197
Gey, K.F., see Asmis, R. 691(1997)59
Godbillon, J., see Sandrenan, N. 691(1997)145
Gundersen, Th.E., Lundanes, E. and Blomhoff, R.
Quantitative high-performance liquid chromatographic determination of retinoids in human serum using on-line solid-phase extraction and column switching. Determination of 9-*cis*-retinoic acid, 13-*cis*-retinoic acid, all-*trans*-retinoic acid, 4-oxo-all-*trans*-retinoic acid and 4-oxo-13-*cis*-retinoic acid 691(1997)43
Halaris, A.E., see Feng, Y. 691(1997)277
Hampl, R., see Hill, M. 691(1997)187
Hanocq, M., see Tassin, J.P. 691(1997)449
Haupt, K., see Legallais, C. 691(1997)33
Heimbecher, S., Lee, Y.-C., Tabibi, S.E. and Yalkowsky, S.H.
Derivatization and high-performance liquid chromatographic analysis of pentaazapentacosane pentahydrochloride 691(1997)173
Hemminki, K., see Koskinen, M. 691(1997)155
Henion, J., see Cai, J. 691(1997)357
Hennekes, M.W.T., see Teerlink, T. 691(1997)269
Hibiya, M., Teradaira, R., Shimpo, K., Matsui, T., Sugimoto, T. and Nagatsu, T.
Interference of a methotrexate derivative with urinary oncopterin [N^2-(3-aminopropyl)biopterin] measurement by high-performance liquid chromatography with fluorimetric detection 691(1997)223

Hill, M., Lapčík, O. and Hampl, R.
Evaluation and separation of steroid–bovine serum albumin conjugates by high-performance liquid chromatography 691(1997)187
Hubbard, J.W., see Yan, J.-h. 691(1997)131
Huwyler, J., see Drewe, J. 691(1997)105
Imamura, T., see Tokunaga, H. 691(1997)203
Ishida, Y., see Takeuchi, A. 691(1997)313
Jackson, J., see Kijak, P.J. 691(1997)377
Jelk, J., see Asmis, R. 691(1997)59
Jitsufuchi, N., see Tokunaga, H. 691(1997)203
Jönsson, B.A.G., see Lindh, C.H. 691(1997)331
Karnes, H.T., see Ramaiya, A. 691(1997)119
Kenndler, E., see Morand, M. 691(1997)192
Kennedy, D.G., see McCracken, R.J. 691(1997)87
Kijak, P.J., Jackson, J. and Shaikh, B.
Determination of gentamicin in bovine milk using liquid chromatography with post-column derivatization and fluorescence detection 691(1997)377
Kimura, A., see Murai, T. 691(1997)13
Kitamura, R., see Matsushima, E. 691(1997)95
Kobayashi, T., see Takeuchi, A. 691(1997)313
Koskinen, M., Rajaniemi, H. and Hemminki, K.
Analysis of tamoxifen-induced DNA adducts by ^{32}P-postlabelling assay using different chromatographic techniques 691(1997)155
Kudo, K., see Tokunaga, H. 691(1997)203
Kurosawa, T., see Murai, T. 691(1997)13
Küsters, E., see Drewe, J. 691(1997)105
Lachno, D.R., see Brooks, S.A. 691(1997)383
Lamprecht, G.
Rapid method for the isolation of hydroxylysylpyridinoline and lysylpyridinoline from concentrated bone hydrolysates by liquid chromatographic techniques 691(1997)297
Lapčík, O., see Hill, M. 691(1997)187
Lasch, J., see Zellmer, S. 691(1997)321
Lee, H.G. and Desiderio, D.M.
Optimization of the capillary zone electrophoresis loading limit and resolution of proteins, using triethylamine, ammonium formate and acidic pH 691(1997)67
Lee, Y.-C., see Heimbecher, S. 691(1997)173
Legallais, C., Anspach, F.B., Bueno, S.M.A., Haupt, K. and Vijayalakshmi, M.A.
Strategies for the depyrogenation of contaminated immunoglobulin G solutions by histidine-immobilized hollow fiber membrane 691(1997)33
Legayada, E., see El-Yazigi, A. 691(1997)457
Li, R.C., see Chiu, F.C.K. 691(1997)473
Lindh, C.H. and Jönsson, B.A.G.
Determination of hexahydrophthalic acid and methylhexahydrophthalic acid in plasma after derivatisation with pentafluorobenzyl bromide using gas chromatography and mass spectrometric detection 691(1997)331
Lo Greco, P., see Londero, D. 691(1997)139
Locher, M., see Paris, S. 691(1997)463
Londero, D. and Lo Greco, P.
New micromethod for the determination of lamotrigine in human plasma by high-performance liquid chromatography 691(1997)139
Lundanes, E., see Gundersen, Th.E. 691(1997)43
Mahara, R., see Murai, T. 691(1997)13
Maher, T.J., see McCabe, D.R. 691(1997)23
Manabe, S., Gao, Q., Yuan, J., Takahashi, T. and Ueki, A.
Determination of isatin in urine and plasma by high-performance liquid chromatography 691(1997)197
Manouilov, K.K., see Chen, H. 691(1997)425
Marquer, C. and Bressolle, F.
High-performance liquid chromatographic determination of the conjugate metabolites of moxisylyte in human plasma and urine 691(1997)389
Masuda, S., see Takeuchi, A. 691(1997)313
Matsuda, I., see Takeuchi, A. 691(1997)313
Matsui, T., see Hibiya, M. 691(1997)223
Matsushima, E., Yoshida, K., Kitamura, R. and Yoshida, K.-i.
Determination of S-1 (combined drug of tegafur, 5-chloro-2,4-dihydroxypyridine and potassium oxonate) and 5-fluorouracil in human plasma and urine using high-performance liquid chromatography and gas chromatography–negative ion chemical ionization mass spectrometry 691(1997)95
Matuszewski, B.K., see Chavez-Eng, C.M. 691(1997)77
McCabe, D.R., Maher, T.J. and Acworth, I.N.
Improved method for the estimation of hydroxyl free radical levels in vivo based on liquid chromatography with electrochemical detection 691(1997)23
McClure, H.M., see Chen, H. 691(1997)425
McCracken, R.J. and Kennedy, D.G.
Determination of the furazolidone metabolite, 3-amino-2-oxazolidinone, in porcine tissues using liquid chromatography–thermospray mass spectrometry and the occurrence of residues in pigs produced in Northern Ireland 691(1997)87
McKay, G., see Yan, J.-h. 691(1997)131
Midha, K.K., see Yan, J.-h. 691(1997)131
Möller, L., see Zeisig, M. 691(1997)341
Morand, M., Blaas, D. and Kenndler, E.
Reduction of wall adsorption in capillary zone electrophoresis of a basic single-chain antibody fragment by a cationic polymeric buffer additive 691(1997)192
Morino, M., see Yoshioka, E. 691(1997)371
Mulder, C., see Teerlink, T. 691(1997)269
Murai, T., Mahara, R., Kurosawa, T., Kimura, A. and Tohma, M.
Determination of fetal bile acids in biological fluids from neonates by gas chromatography–negative ion chemical ionization mass spectrometry 691(1997)13
Nagatsu, T., see Hibiya, M. 691(1997)223
Naidu, M.U.R., see Ravi, S.K. 691(1997)208
Netter, C., see Sandrenan, N. 691(1997)145
Nishiyama, S., see Takeuchi, A. 691(1997)313
Nowitzki-Grimm, S., see Schieber, A. 691(1997)1
Obermeyer, B.D., see Brooks, S.A. 691(1997)383
Ohta, H., Seto, Y. and Tsunoda, N.
Determination of *Aconitum* alkaloids in blood and urine samples. I. High-performance liquid chromatographic separation, solid-phase extraction and mass spectrometric confirmation 691(1997)351
Ohtsuka, Y., see Tokunaga, H. 691(1997)203
Okano, T., see Takeuchi, A. 691(1997)313
Paibir, S.G. and Soine, W.H.
High-performance liquid chromatographic analysis of phenobarbital and phenobarbital metabolites in human urine 691(1997)111

Paris, S., Blaschke, G., Locher, M., Borbe, H.O. and Engel, J.
Investigation of the stereoselective in vitro metabolism of the chiral antiasthmatic/antiallergic drug flezelastine by high-performance liquid chromatography and capillary zone electrophoresis 691(1997)463
Piletz, J.E., see Feng, Y. 691(1997)277
Rademaker, R.V., see Saris, L.A. 691(1997)409
Rajaniemi, H., see Koskinen, M. 691(1997)155
Ramaiya, A. and Karnes, H.T.
Simultaneous measurement of serotonin and paroxetine in rat brain microdialysate by a single-pump column-switching technique 691(1997)119
Ramzan, I., see Yeganeh, M.H. 691(1997)478
Rani, P.U., see Ravi, S.K. 691(1997)208
Rao, T.R.K., see Ravi, S.K. 691(1997)208
Ravi, S.K., Naidu, M.U.R., Sekhar, E.C., Rao, T.R.K., Shobha, J.C., Rani, P.U. and Surya, K.J.
Rapid and selective analysis of secnidazole in human plasma using high-performance liquid chromatography with ultraviolet detection 691(1997)208
Raymond, K., see Chiu, F.C.K. 691(1997)473
Rooyakkers, D.R., see Van Eijk, H.M.H. 691(1997)287
Rufer, S., see Drewe, J. 691(1997)105
Rupp-Classen, M., see Schieber, A. 691(1997)1
Sandrenan, N., Sioufi, A., Godbillon, J., Netter, C., Donker, M. and van Valkenburg, C.
Determination of artemether and its metabolite, dihydroartemisinin, in plasma by high-performance liquid chromatography and electrochemical detection in the reductive mode 691(1997)145
Saris, L.A., Brekelmans, G.J.F., van der Linden, G.J., Rademaker, R.V. and Edelbroek, P.M.
High-performance liquid chromatographic determination of carbamazepine and metabolites in human hair 691(1997)409
Schieber, A., Brückner, H., Rupp-Classen, M., Specht, W., Nowitzki-Grimm, S. and Classen, H.-G.
Evaluation of D-amino acid levels in rat by gas chromatography–selected ion monitoring mass spectrometry: no evidence for subacute toxicity of orally fed D-proline and D-aspartic acid 691(1997)1
Schinazi, R.F., see Chen, H. 691(1997)425
Sekhar, E.C., see Ravi, S.K. 691(1997)208
Sekimoto, H., see Takeuchi, A. 691(1997)313
Selley, M.L.
Determination of the lipid peroxidation product (*E*)-4-hydroxy-2-nonenal in clinical samples by gas chromatography–negative-ion chemical ionisation mass spectrometry of the *O*-pentafluorobenzyl oxime 691(1997)263
Seto, Y., see Ohta, H. 691(1997)351
Shaikh, B., see Kijak, P.J. 691(1997)377
Shimpo, K., see Hibiya, M. 691(1997)223
Shinkai, H., see Yoshioka, E. 691(1997)371
Shobha, J.C., see Ravi, S.K. 691(1997)208
Siluveru, M. and Stewart, J.T.
Enantioselective determination of *S*-(+)- and *R*-(−)-ondansetron in human serum using derivatized cyclodextrin-modified capillary electrophoresis and solid-phase extraction 691(1997)217
Singer, M., see Timm, U. 691(1997)397
Sioufi, A., see Sandrenan, N. 691(1997)145
Soeters, P.B., see Van Eijk, H.M.H. 691(1997)287
Soine, W.H., see Paibir, S.G. 691(1997)111
Sommadossi, J.-P., see Zhou, X.-J. 691(1997)417
Specht, W., see Schieber, A. 691(1997)1
Stankovic, C.J., see Surendran, N. 691(1997)305
Steiner, B., see Timm, U. 691(1997)397
Stewart, B.H., see Surendran, N. 691(1997)305
Stewart, J.T., see Siluveru, M. 691(1997)217
Stopher, D.A. and Gage, R.
Determination of a new antifungal agent, voriconazole, by multidimensional high-performance liquid chromatography with direct plasma injection onto a size-exclusion column 691(1997)441
Sugimoto, T., see Hibiya, M. 691(1997)223
Surendran, N., Stankovic, C.J. and Stewart, B.H.
Development and validation of an ion-pair reversed-phase high-performance liquid chromatographic method for the separation of phosphonodipeptides 691(1997)305
Surya, K.J., see Ravi, S.K. 691(1997)208
Suskin, M., see Berger, B.J. 691(1997)433
Tabibi, S.E., see Heimbecher, S. 691(1997)173
Takahashi, T., see Manabe, S. 691(1997)197
Takeuchi, A., Ishida, Y., Sekimoto, H., Masuda, S., Okano, T., Nishiyama, S., Matsuda, I. and Kobayashi, T.
Simplified method for the determination of 25-hydroxy and 1α,25-dihydroxy metabolites of vitamins D_2 and D_3 in human plasma. Application to nutritional studies 691(1997)313
Tanaka, H., see Yoshioka, E. 691(1997)371
Tassin, J.P., Dubois, J., Atassi, G. and Hanocq, M.
Simultaneous determination of cytotoxic (adriamycin, vincristine) and modulator of resistance (verapamil, S 9788) drugs in human cells by high-performance liquid chromatography and ultraviolet detection 691(1997)449
Teerlink, T., Hennekes, M.W.T., Mulder, C. and Brulez, H.F.H.
Determination of dimethylamine in biological samples by high-performance liquid chromatography 691(1997)269
Teradaira, R., see Hibiya, M. 691(1997)223
Timm, U., Zumbrunnen, R., Erdin, R., Singer, M. and Steiner, B.
Oral platelet aggregation inhibitor Ro 48-3657: Determination of the active metabolite and its prodrug in plasma and urine by high-performance liquid chromatography using automated column switching 691(1997)397
Tohma, M., see Murai, T. 691(1997)13
Tokunaga, H., Kudo, K., Jitsufuchi, N., Ohtsuka, Y. and Imamura, T.
Sensitive determination of sulpiride in human plasma by high- performance liquid chromatography 691(1997)203
Tsunoda, N., see Ohta, H. 691(1997)351
Ueki, A., see Manabe, S. 691(1997)197
Ulrich, P., see Berger, B.J. 691(1997)433
Urakawa, A., see Azuma, R. 691(1997)179
Van der Linden, G.J., see Saris, L.A. 691(1997)409
Van Eijk, H.M.H., Rooyakkers, D.R., Wagenmakers, A.J.M., Soeters, P.B. and Deutz, N.E.P.
Isolation and quantitation of isotopically labeled amino acids from biological samples 691(1997)287
Van Valkenburg, C., see Sandrenan, N. 691(1997)145
Vijayalakshmi, M.A., see Legallais, C. 691(1997)33
Wagenmakers, A.J.M., see Van Eijk, H.M.H. 691(1997)287
Warner, D.L. and Burke, T.G.
Simple and versatile high-performance liquid chromatographic method for the simultaneous quantitation of the lactone and carboxylate forms of camptothecin anticancer drugs 691(1997)161

Yalkowsky, S.H., see Heimbecher, S. 691(1997)173
Yan, J.-h., Hubbard, J.W., McKay, G. and Midha, K.K.
Stereoselective and simultaneous measurement of *cis*- and *trans*-isomers of doxepin and N-desmethyldoxepin in plasma or urine by high-performance liquid chromatography 691(1997)131
Yeganeh, M.H. and Ramzan, I.
Determination of propofol in rat whole blood and plasma by high-performance liquid chromatography 691(1997)478
Yoshida, K., see Matsushima, E. 691(1997)95
Yoshida, K.-i., see Matsushima, E. 691(1997)95
Yoshioka, E., Morino, M., Tanaka, H. and Shinkai, H.
Manual and automatic extraction and high-performance liquid chromatographic determination of a spicamycin derivative, KRN5500, in rat plasma 691(1997)371
Yuan, J., see Manabe, S. 691(1997)197
Zeisig, M. and Möller, L.
^{32}P-Postlabeling high-performance liquid chromatographic improvements to characterize DNA adduct stereoisomers from benzo[*a*]pyrene and benzo[*c*]phenanthrene, and to separate DNA adducts from 7,12-dimethylbenz[*a*]anthracene 691(1997)341
Zellmer, S. and Lasch, J.
Individual variation of human plantar stratum corneum lipids, determined by automated multiple development of high-performance thin-layer chromatography plates 691(1997)321
Zhang, J.N., see Chiu, F.C.K. 691(1997)473
Zhou, X.-J. and Sommadossi, J.-P.
Rapid quantitation of (−)-2′-deoxy-3′-thiacytidine in human serum by high-performance liquid chromatography with ultraviolet detection 691(1997)417
Zumbrunnen, R., see Timm, U. 691(1997)397

Compound Index Vol. 691

***Aconitum* alkaloids**
Determination of *Aconitum* alkaloids in blood and urine samples. I. High-performance liquid chromatographic separation, solid-phase extraction and mass spectrometric confirmation 691(1997)351

Adriamycin
Simultaneous determination of cytotoxic (adriamycin, vincristine) and modulator of resistance (verapamil, S 9788) drugs in human cells by high-performance liquid chromatography and ultraviolet detection 691(1997)449

Agmatine
Determination of agmatine in brain and plasma using high-performance liquid chromatography with fluorescence detection 691(1997)277

β-Agonists
Quantitative multi-residue determination of β-agonists in bovine urine using on-line immunoaffinity extraction–coupled column packed capillary liquid chromatography–tandem mass spectrometry 691(1997)357

Alkaloids
Determination of *Aconitum* alkaloids in blood and urine samples. I. High-performance liquid chromatographic separation, solid-phase extraction and mass spectrometric confirmation 691(1997)351

Amino acids
Isolation and quantitation of isotopically labeled amino acids from biological samples 691(1997)287

Rapid method for the isolation of hydroxylysylpyridinoline and lysylpyridinoline from concentrated bone hydrolysates by liquid chromatographic techniques 691(1997)297

D-Amino acids
Evaluation of D-amino acid levels in rat by gas chromatography–selected ion monitoring mass spectrometry: no evidence for subacute toxicity of orally fed D-proline and D-aspartic acid 691(1997)1

2-Amino-6-chloropurine dioxolane
High-performance liquid chromatographic determination of (−)-β-D-2-aminopurine dioxolane and (−)-β-D-2-amino-6-chloropurine dioxolane, and their metabolite (−)-β-D-dioxolane guanine in monkey serum, urine and cerebrospinal fluid 691(1997)425

2-Amino-2-oxazolidinone
Determination of the furazolidone metabolite, 3-amino-2-oxazolidinone, in porcine tissues using liquid chromatography–thermospray mass spectrometry and the occurrence of residues in pigs produced in Northern Ireland 691(1997)87

2-Aminopurine dioxolane
High-performance liquid chromatographic determination of (−)-β-D-2-aminopurine dioxolane and (−)-β-D-2-amino-6-chloropurine dioxolane, and their metabolite (−)-β-D-dioxolane guanine in monkey serum, urine and cerebrospinal fluid 691(1997)425

Ammonium formate
Optimization of the capillary zone electrophoresis loading limit and resolution of proteins, using triethylamine, ammonium formate and acidic pH 691(1997)67

Arginine
Determination of agmatine in brain and plasma using high-performance liquid chromatography with fluorescence detection 691(1997)277

Artemether
Determination of artemether and its metabolite, dihydroartemisinin, in plasma by high-performance liquid chromatography and electrochemical detection in the reductive mode 691(1997)145

Arylene bis(methylketone) compounds
Studies on the pharmacological properties of novel arylene bis(methylketone) compounds using solid-phase extraction and high-performance liquid chromatography 691(1997)433

D-Aspartic acid
Evaluation of D-amino acid levels in rat by gas chromatography–selected ion monitoring mass spectrometry: no evidence for subacute toxicity of orally fed D-proline and D-aspartic acid 691(1997)1

Atenolol
Efficient assay for the determination of atenolol in human plasma and urine by high-performance liquid chromatography with fluorescence detection 691(1997)473

Benzo[*c*]phenanthrene
^{32}P-Postlabeling high-performance liquid chromatographic improvements to characterize DNA adduct stereoisomers from benzo[*a*]pyrene and benzo[*c*]phenanthrene, and to separate DNA adducts from 7,12-dimethylbenz[*a*]anthracene 691(1997)341

Benzo[*a*]pyrene
^{32}P-Postlabeling high-performance liquid chromatographic improvements to characterize DNA adduct stereoisomers from benzo[*a*]pyrene and benzo[*c*]phenanthrene, and to separate DNA adducts from 7,12-dimethylbenz[*a*]**anthracene** 691(1997)341

Benzoquinolinones
High-performance liquid chromatographic determination of the enantiomers of the benzoquinolinone LY191704, a human type 1 5α-reductase inhibitor, in plasma 691(1997)212

Bile acids
Determination of fetal bile acids in biological fluids from neonates by gas chromatography–negative ion chemical ionization mass spectrometry 691(1997)13

Bovine serum albumin
Evaluation and separation of steroid–bovine serum albumin conjugates by high-performance liquid chromatography 691(1997)187

Bupivacaine
High-performance liquid chromatographic method for an automated determination of local anaesthetics in human plasma 691(1997)105

Camptothecin
Simple and versatile high-performance liquid chromatographic method for the simultaneous quantitation of the lactone and carboxylate forms of camptothecin anticancer drugs 691(1997)161

Carbamazepine
High-performance liquid chromatographic determination of carbamazepine and metabolites in human hair 691(1997)409

5-Chloro-2,4-dihydroxypyridine
Determination of S-1 (combined drug of tegafur, 5-chloro-2,4-dihydroxypyridine and potassium oxonate) and 5-fluorouracil in human plasma and urine using high-performance liquid chromatography and gas chromatography–negative ion chemical ionization mass spectrometry 691(1997)95

Cholesterol
Concurrent quantification of cellular cholesterol, cholesteryl esters and triglycerides in small biological samples 691(1997)59

Cholesteryl esters
Concurrent quantification of cellular cholesterol, cholesteryl esters and triglycerides in small biological samples 691(1997)59

Clenbuterol
Quantitative multi-residue determination of β-agonists in bovine urine using on-line immunoaffinity extraction–coupled column packed capillary liquid chromatography–tandem mass spectrometry 691(1997)357

Dansylation
Derivatization and high-performance liquid chromatographic analysis of pentaazapentacosane pentahydrochloride 691(1997)173

Daunorubicin
Simultaneous determination of cytotoxic (adriamycin, vincristine) and modulator of resistance (verapamil, S 9788) drugs in human cells by high-performance liquid chromatography and ultraviolet detection 691(1997)449

Deacetylmoxisylyte sulphoconjugate
High-performance liquid chromatographic determination of the conjugate metabolites of moxisylyte in human plasma and urine 691(1997)389

Demethyl-TAS-103
Simultaneous determination of a novel anticancer drug, TAS-103, and its N-demethylated metabolite in monkey plasma by high-performance liquid chromatography using solid-phase extraction 691(1997)179

(−)-2′-Deoxy-3′-thiacytidine
Rapid quantitation of (−)-2′-deoxy-3′-thiacytidine in human serum by high-performance liquid chromatography with ultraviolet detection 691(1997)417

N-Desmethyldoxepin
Stereoselective and simultaneous measurement of *cis*- and *trans*-isomers of doxepin and N-desmethyldoxepin in plasma or urine by high-performance liquid chromatography 691(1997)131

Dihydroartemisinin
Determination of artemether and its metabolite, dihydroartemisinin, in plasma by high-performance liquid chromatography and electrochemical detection in the reductive mode 691(1997)145

Dihydroxybenzoic acid
Improved method for the estimation of hydroxyl free radical levels in vivo based on liquid chromatography with electrochemical detection 691(1997)23

Dimethylamine
Determination of dimethylamine in biological samples by high-performance liquid chromatography 691(1997)269

6-{[2-(Dimethylamino)ethyl]amino}-3-hydroxy-7*H*-indeno[2,1-*c*]quinolin-7-one dihydrochloride
Simultaneous determination of a novel anticancer drug, TAS-103, and its N-demethylated metabolite in monkey plasma by high-performance liquid chromatography using solid-phase extraction 691(1997)179

7,12-Dimethylbenz[*a*]anthracene
^{32}P-Postlabeling high-performance liquid chromatographic improvements to characterize DNA adduct stereoisomers from benzo[*a*]pyrene and benzo[*c*]phenanthrene, and to separate DNA adducts from 7,12-dimethylbenz[*a*]anthracene 691(1997)341

Dioxolane guanine
High-performance liquid chromatographic determination of (−)-β-D-2-aminopurine dioxolane and (−)-β-D-2-amino-6-chloropurine dioxolane, and their metabolite (−)-β-D-dioxolane guanine in monkey serum, urine and cerebrospinal fluid 691(1997)425

DNA
^{32}P-Postlabeling high-performance liquid chromatographic improvements to characterize DNA adduct stereoisomers from benzo[*a*]pyrene and benzo[*c*]phenanthrene, and to separate DNA adducts from 7,12-dimethylbenz[*a*]anthracene 691(1997)341

DNA adducts
Analysis of tamoxifen-induced DNA adducts by ^{32}P-postlabelling assay using different chromatographic techniques 691(1997)155

Dopamine D_4 receptor antagonists
Picogram determination of a novel dopamine D_4 receptor antagonist in human plasma and urine by liquid chromatography with atmospheric pressure chemical ionization tandem mass spectrometry 691(1997)77

Doxepin
Stereoselective and simultaneous measurement of *cis*- and *trans*-isomers of doxepin and N-desmethyldoxepin in plasma or urine by high-performance liquid chromatography 691(1997)131

Endotoxins
Strategies for the depyrogenation of contaminated immunoglobulin G solutions by histidine-immobilized hollow fiber membrane 691(1997)33

Ergolines
Automated high-performance liquid chromatographic method for the analysis of two novel ergoline compounds in human plasma 691(1997)383

Etidocaine
High-performance liquid chromatographic method for an automated determination of local anaesthetics in human plasma 691(1997)105

Fatty acids
Concurrent quantification of cellular cholesterol, cholesteryl esters and triglycerides in small biological samples 691(1997)59

Individual variation of human plantar stratum corneum lipids, determined by automated multiple development of high-performance thin-layer chromatography plates 691(1997)321

Flezelastine
Investigation of the stereoselective in vitro metabolism of the chiral antiasthmatic/antiallergic drug flezelastine by high-performance liquid chromatography and capillary zone electrophoresis 691(1997)463

5-Fluorouracil
Determination of S-1 (combined drug of tegafur, 5-chloro-2,4-dihydroxypyridine and potassium oxonate) and 5-fluorouracil in human plasma and urine using high-performance liquid chromatography and gas chromatography–negative ion chemical ionization mass spectrometry 691(1997)95

Furazolidone
Determination of the furazolidone metabolite, 3-amino-2-oxazolidinone, in porcine tissues using liquid chromatography–thermospray mass spectrometry and the occurrence of residues in pigs produced in Northern Ireland 691(1997)87

Gentamicin
Determination of gentamicin in bovine milk using liquid chromatography with post-column derivatization and fluorescence detection 691(1997)377

Hexahydrophthalic acid
Determination of hexahydrophthalic acid and methylhexahydrophthalic acid in plasma after derivatisation with pentafluorobenzyl bromide using gas chromatography and mass spectrometric detection 691(1997)331

Histidine
Strategies for the depyrogenation of contaminated immunoglobulin G solutions by histidine-immobilized hollow fiber membrane 691(1997)33

Hydroxyl free radicals
Improved method for the estimation of hydroxyl free radical levels in vivo based on liquid chromatography with electrochemical detection 691(1997)23

Hydroxylysylpyridinoline
Rapid method for the isolation of hydroxylysylpyridinoline and lysylpyridinoline from concentrated bone hydrolysates by liquid chromatographic techniques 691(1997)297

(*E*)-4-Hydroxy-2-nonenol
Determination of the lipid peroxidation product (*E*)-4-hydroxy-2-nonenal in clinical samples by gas chromatography–negative-ion chemical ionisation mass spectrometry of the *O*-pentafluorobenzyl oxime 691(1997)263

***p*-Hydroxyphenobarbital**
High-performance liquid chromatographic analysis of phenobarbital and phenobarbital metabolites in human urine 691(1997)111

Imidazoline receptors
Determination of agmatine in brain and plasma using high-performance liquid chromatography with fluorescence detection 691(1997)277

Immunoglobulin G
Strategies for the depyrogenation of contaminated immunoglobulin G solutions by histidine-immobilized hollow fiber membrane 691(1997)33

Indole-2,3-dione
Determination of isatin in urine and plasma by high-performance liquid chromatography 691(1997)197

Isatin
Determination of isatin in urine and plasma by high-performance liquid chromatography 691(1997)197

KRN5500
Manual and automatic extraction and high-performance liquid chromatographic determination of a spicamycin derivative, KRN5500, in rat plasma 691(1997)371

α-Lactalbumin
Optimization of the capillary zone electrophoresis loading limit and resolution of proteins, using triethylamine, ammonium formate and acidic pH 691(1997)67

Lamotrigine
New micromethod for the determination of lamotrigine in human plasma by high-performance liquid chromatography 691(1997)139

Lipids
Determination of the lipid peroxidation product (*E*)-4-hydroxy-2-nonenal in clinical samples by gas chromatography–negative-ion chemical ionisation mass spectrometry of the *O*-pentafluorobenzyl oxime 691(1997)263

Individual variation of human plantar stratum corneum lipids, determined by automated multiple development of high-performance thin-layer chromatography plates 691(1997)321

LY191704
High-performance liquid chromatographic determination of the enantiomers of the benzoquinolinone LY191704, a human type 1 5α-reductase inhibitor, in plasma 691(1997)212

LY300502
High-performance liquid chromatographic determination of the enantiomers of the benzoquinolinone LY191704, a human type 1 5α-reductase inhibitor, in plasma 691(1997)212

Lysozyme
Optimization of the capillary zone electrophoresis loading limit and resolution of proteins, using triethylamine, ammonium formate and acidic pH 691(1997)67

Lysylpyridinoline
Rapid method for the isolation of hydroxylysylpyridinoline and lysylpyridinoline from concentrated bone hydrolysates by liquid chromatographic techniques 691(1997)297

Mabuterol
Quantitative multi-residue determination of β-agonists in bovine urine using on-line immunoaffinity extraction–coupled column packed capillary liquid chromatography–tandem mass spectrometry 691(1997)357

Mapenterol
Quantitative multi-residue determination of β-agonists in bovine urine using on-line immunoaffinity extraction–coupled column packed capillary liquid chromatography–tandem mass spectrometry 691(1997)357

Methotrexate
Interference of a methotrexate derivative with urinary oncopterin [N^2-(3-aminopropyl)biopterin] measurement by high-performance liquid chromatography with fluorimetric detection 691(1997)223

Methylclenbuterol
Quantitative multi-residue determination of β-agonists in bovine urine using on-line immunoaffinity extraction–coupled column packed capillary liquid chromatography–tandem mass spectrometry 691(1997)357

Methylhexahydrophthalic acid
Determination of hexahydrophthalic acid and methylhexahydrophthalic acid in plasma after derivatisation with pentafluorobenzyl bromide using gas chromatography and mass spectrometric detection 691(1997)331

Monomethyldeacetylmoxisylyte sulphoconjugate
High-performance liquid chromatographic determination of the conjugate metabolites of moxisylyte in human plasma and urine 691(1997)389

Moxisylyte
High-performance liquid chromatographic determination of the conjugate metabolites of moxisylyte in human plasma and urine 691(1997)389

Myoglobin
Optimization of the capillary zone electrophoresis loading limit and resolution of proteins, using triethylamine, ammonium formate and acidic pH 691(1997)67

Non-steroidal anti-inflammatory drugs
Methods of analysis of chiral non-steroidal anti-inflammatory drugs 691(1997)229

N^2-(3-Aminopropyl)biopterin
Interference of a methotrexate derivative with urinary oncopterin [N^2-(3-aminopropyl)biopterin] measurement by high-performance liquid chromatography with fluorimetric detection 691(1997)223

Oncopterin
Interference of a methotrexate derivative with urinary oncopterin [N^2-(3-aminopropyl)biopterin] measurement by high-performance liquid chromatography with fluorimetric detection 691(1997)223

Ondansetron
Enantioselective determination of *S*-(+)- and *R*-(−)-ondansetron in human serum using derivatized cyclodextrin-modified capillary electrophoresis and solid-phase extraction 691(1997)217

Oxoretinoic acid
Quantitative high-performance liquid chromatographic determination of retinoids in human serum using on-line solid-phase extraction and column switching 691(1997)43

Paroxetine
Simultaneous measurement of serotonin and paroxetine in rat brain microdialysate by a single-pump column-switching technique 691(1997)119

Pentaazapentacosane pentahydrochloride
Derivatization and high-performance liquid chromatographic analysis of pentaazapentacosane pentahydrochloride 691(1997)173

Phenobarbital
High-performance liquid chromatographic analysis of phenobarbital and phenobarbital metabolites in human urine 691(1997)111

Phenobarbital N-glucosides
High-performance liquid chromatographic analysis of phenobarbital and phenobarbital metabolites in human urine 691(1997)111

Phenobarbital N-glucuronides
High-performance liquid chromatographic analysis of phenobarbital and phenobarbital metabolites in human urine 691(1997)111

Phosphonodipeptides
Development and validation of an ion-pair reversed-phase high-performance liquid chromatographic method for the separation of phosphonodipeptides 691(1997)305

Polyamines
Derivatization and high-performance liquid chromatographic analysis of pentaazapentacosane pentahydrochloride 691(1997)173

Potassium oxonate
Determination of S-1 (combined drug of tegafur, 5-chloro-2,4-dihydroxypyridine and potassium oxonate) and 5-fluorouracil in human plasma and urine using high-performance liquid chromatography and gas chromatography–negative ion chemical ionization mass spectrometry 691(1997)95

D-Proline
Evaluation of D-amino acid levels in rat by gas chromatography–selected ion monitoring mass spectrometry: no evidence for subacute toxicity of orally fed D-proline and D-aspartic acid 691(1997)1

Propofol
Determination of propofol in rat whole blood and plasma by high-performance liquid chromatography 691(1997)478

Proteins
Optimization of the capillary zone electrophoresis loading limit and resolution of proteins, using triethylamine, ammonium formate and acidic pH 691(1997)67

Reduction of wall adsorption in capillary zone electrophoresis of a basic single-chain antibody fragment by a cationic polymeric buffer additive 691(1997)192

5α-Reductase inhibitor
High-performance liquid chromatographic determination of the enantiomers of the benzoquinolinone LY191704, a human type 1 5α-reductase inhibitor, in plasma 691(1997)212

Retinoic acid
Quantitative high-performance liquid chromatographic determination of retinoids in human serum using on-line solid-phase extraction and column switching 691(1997)43

Retinoids
Quantitative high-performance liquid chromatographic determination of retinoids in human serum using on-line solid-phase extraction and column switching 691(1997)43

Ribonuclease A
Optimization of the capillary zone electrophoresis loading limit and resolution of proteins, using triethylamine, ammonium formate and acidic pH 691(1997)67

Ro 48-3657
Oral platelet aggregation inhibitor Ro 48-3657: Determination of the active metabolite and its prodrug in plasma and urine by high-performance liquid chromatography using automated column switching 691(1997)397

S-1
Determination of S-1 (combined drug of tegafur, 5-chloro-2,4-dihydroxypyridine and potassium oxonate) and 5-fluorouracil in human plasma and urine using high-performance liquid chromatography and gas chromatography–negative ion chemical ionization mass spectrometry 691(1997)95

S 9788
Simultaneous determination of cytotoxic (adriamycin, vincristine) and modulator of resistance (verapamil, S 9788) drugs in human cells by high-performance liquid chromatography and ultraviolet detection 691(1997)449

Salicylic acid
Improved method for the estimation of hydroxyl free radical levels in vivo based on liquid chromatography with electrochemical detection 691(1997)23

Secnidazole
Rapid and selective analysis of secnidazole in human plasma using high-performance liquid chromatography with ultraviolet detection 691(1997)208

Sergolexole
Automated high-performance liquid chromatographic method for the analysis of two novel ergoline compounds in human plasma 691(1997)383

Serotonin
Simultaneous measurement of serotonin and paroxetine in rat brain microdialysate by a single-pump column-switching technique 691(1997)119

Single-chain antibody fragments
Reduction of wall adsorption in capillary zone electrophoresis of a basic single-chain antibody fragment by a cationic polymeric buffer additive 691(1997)192

Spicamycin derivative
Manual and automatic extraction and high-performance liquid chromatographic determination of a spicamycin derivative, KRN5500, in rat plasma 691(1997)371

Steroids
Evaluation and separation of steroid–bovine serum albumin conjugates by high-performance liquid chromatography 691(1997)187

Sulpiride
Sensitive determination of sulpiride in human plasma by high-performance liquid chromatography 691(1997)203

Tamoxifen
Analysis of tamoxifen-induced DNA adducts by ^{32}P-postlabelling assay using different chromatographic techniques 691(1997)155

Direct liquid chromatographic micro-measurement of tamoxifen in plasma of cancer patients 691(1997)457

TAS-103
Simultaneous determination of a novel anticancer drug, TAS-103, and its N-demethylated metabolite in monkey plasma by high-performance liquid chromatography using solid-phase extraction 691(1997)179

Tegafur
Determination of S-1 (combined drug of tegafur, 5-chloro-2,4-dihydroxypyridine and potassium oxonate) and 5-fluorouracil in human plasma and urine using high-performance liquid chromatography and gas chromatography–negative ion chemical ionization mass spectrometry 691(1997)95

Tolubuterol
Quantitative multi-residue determination of β-agonists in bovine urine using on-line immunoaffinity extraction–coupled column packed capillary liquid chromatography–tandem mass spectrometry 691(1997)357

trans-10,11-Dihydro-10,11-dihydroxycarbamazepine
High-performance liquid chromatographic determination of carbamazepine and metabolites in human hair 691(1997)409

Triethylamine
Optimization of the capillary zone electrophoresis loading limit and resolution of proteins, using triethylamine, ammonium formate and acidic pH 691(1997)67

Triglycerides
Concurrent quantification of cellular cholesterol, cholesteryl esters and triglycerides in small biological samples 691(1997)59

Trypsinogen
Optimization of the capillary zone electrophoresis loading limit and resolution of proteins, using triethylamine, ammonium formate and acidic pH 691(1997)67

Verapamil
Simultaneous determination of cytotoxic (adriamycin, vincristine) and modulator of resistance (verapamil, S 9788) drugs in human cells by high-performance liquid chromatography and ultraviolet detection 691(1997)449

Vincristine
Simultaneous determination of cytotoxic (adriamycin, vincristine) and modulator of resistance (verapamil, S 9788) drugs in human cells by high-performance liquid chromatography and ultraviolet detection 691(1997)449

Voriconazole
Determination of a new antifungal agent, voriconazole, by multidimensional high-performance liquid chromatography with direct plasma injection onto a size-exclusion column 691(1997)441